



Chmelářský institut s. r. o.

HOP PROTECTION AGAINST ALFALFA SNOUT BEETLE (*Otiorhynchus ligustici* L.) WITH THE HELP OF METEOROLOGICAL DATA IN BOHEMIAN AND MORAVIAN HOP GARDENS

Josef Vostřel, Ivo Klopal, Tomáš Kudrna

Hop Research Institute, Co., Ltd., Kadaňská 2525, Žatec, Czech Republic, e-mail: j.vostrel@telecom.cz



Figure 1: Adult alfalfa snout beetle (*Otiorhynchus ligustici* L.) sitting on a young hop shoot.

The family Curculionidae contains more than 40,000 species, all of which have a head that is extended into a long snout or rostrum. The jaws are on the end of the rostrum and the antennae are implanted in the middle (Malais & Ravensberg, 1992). Some of them cause damage on roots of various plants. Numerous species of root weevils, *Otiorhynchus* spp. infest hop (*Humulus lupulus* L.). Whereas black vine weevil (*O. sulcatus* F.) is dominant species infesting hop in Washington and Oregon hop yards (Mahaffee et al., 2009), alfalfa snout beetle (*Otiorhynchus ligustici* L.) is the most important pest from this family in Czech and Moravian hop gardens (Vostřel, 1999).

The wingless beetles are about 9-13 cm long of black to black-brown colour. They usually shelter under clods of earth during the day where they are difficult to see since their colour blends with the soil. They are active in the spring when they lay eggs in the soil from which the legless larvae develop and feed on the roots before pupating (Neve, 1991). Severe larval infestation can significantly shorten the life of a hop yard. The larvae of this beetle are not easily controlled chemically because they live in the soil in hop crowns. In trials carried out to try to control larvae in a nontraditional way good results were obtained neither from the pesticides applied in the form of watering in the autumn nor from granular insecticides applied onto soil surface to the hop plants and by injection into the rootstock (Vostřel, 1998). The only efficient way to control *O. ligustici* is to apply an efficient insecticide in the spring when newly born adults emerge from soil and damage young hop shoots. In the past treatment was recommended when 100 beetles per 100 plants were found out (Petřík & Štys, 1988). Nevertheless, population density of weevils in Czech a Moravian hop gardens has decreased in comparison with the recent years and therefore it has been necessary to overestimate the economic threshold, which is now 10 beetles per 100 plants. Besides counting beetles at the soil surface on hop shoots and under clods of earth, soil temperature in the depth of 50 cm is measured with the help of the soil probe. It was found out that beetles began to emerge from soil when temperature in 50 cm reaches the value of 8 °C. Nevertheless, 13-15 °C in the above-mentioned depth is necessary for mass emergence of beetles. It usually happens in the second half of April, which is also the recommended time for treatment of young shoots to control this pest and to prevent oviposition by females. The occurrence of alfalfa snout beetle (*Otiorhynchus ligustici* L.) was monitored in the hop gardens within Moravian (Tršice) hop region in 2009-2011 (Table 1-3, Figure 1-2).

REFERENCES:

- Mahaffee, W.F., Pethybridge, S.J., Gent, D.H. (2009):** Compendium of hop diseases and pests. APS (The American Phytopathological Society) press, S. Paul, Minnesota: 93 pp.
Malais M., Ravensberg W.J. (1992): Knowing and Recognizing. In: The biology of glasshouse pests and their natural enemies. Koppert B.V., Berkel en Rodenrijs, the Netherlands, 87-89.
Neve, R.A. (1991): Hops. Chapman and Hall, London: 266 pp.
Petřík, Z., Štys, Z. (1988): Prognosis of diseases and pests in integrated control on hops in Czechoslovakia. In IOBC WPBS Bull. XI/5, East Malling, England: 49-55.
Vostřel, J. (1998): Using some non-traditional methods in the control of hop pests. (In Czech). Rostl. Vyroba, 34: 771-777.
Vostřel, J. (1999): Hop protection against pests and diseases in Czech Republic (up-to-date information). In Proc. Scient. Comm. I.H.G.C., Pulawy, Poland: 120-122.

TABLE 1: The occurrence of alfalfa snout beetle (*Otiorhynchus ligustici* L.) in the hop gardens within Moravian (Tršice) hop-growing region in 2009



LOCALITY	DATE OF EMERGENCY	ALTITUDE (m)	SLOPE	ORIENTATION OF THE LOCALITY	DRAINAGE	YEAR OF PLANTING
Lipany	09.04.	266	2.7°	E 77%	Yes	1995
				NE 23%		
Kokory	05.04.	253	3.2°	W 24%	No	1978
				SW 76%		
Prusy	11.04.	222	2.0°	NE 39%	Yes	1999
				N 53%		
				Plain 8%		
Čechovice	13.04.	256	2.2°	NE 27%	No	1998
				N 9%		
				NW 11%		
				W 33%		
				SW 8%		
Plain 12%						

TABLE 2: The occurrence of alfalfa snout beetle (*Otiorhynchus ligustici* L.) in the hop gardens within Moravian (Tršice) hop-growing region in 2010

LOCALITY	DATE OF EMERGENCY	ALTITUDE (m)	SLOPE	ORIENTATION OF THE LOCALITY	YEAR OF PLANTING	PRUNING
Kokory	19.04.	253 m	3.2°	W 24%	1978	08.04.
				SW 76%		
Prosenice	19.04.	226 m	1.2°	S 24%	1995	31.03.
				SE 16%		
				Plain 60%		
Lazničky	20.04.	283 m	1.1°	SW 62%	1997	16.04.
				NW 1%		
				W 2%		
				Plain 35%		
Lipník nad Bečvou	20.04.	229 m	0.9°	E 1%	2001	04.04.
				W 1%		
				SW 10%		
				S 23%		
				SE 3%		
Plain 62%						
Čechovice	23.04.	256 m	2.2°	NE 27%	1998	20.04.
				N 9%		
				NW 11%		
				W 33%		
				SW 8%		
				Plain 12%		

TABLE 3: The occurrence of alfalfa snout beetle (*Otiorhynchus ligustici* L.) in the hop gardens within Moravian (Tršice) hop-growing region in 2011

LOCALITY	DATE OF EMERGENCY	ALTITUDE (m)	SLOPE	ORIENTATION OF THE LOCALITY	YEAR OF PLANTING	PRUNING
Lipník nad Bečvou	06.04.	229 m	0.9°	E 1%	2001	01.04.
				W 1%		
				SW 10%		
				S 23%		
				SE 3%		
Lazničky	07.04.	283 m	1.1°	SW 62%	1997	08.04.
				NW 1%		
				W 2%		
				Plain 35%		
Čechovice	08.04.	256 m	2.2°	NE 27%	1998	10.04.
				N 9%		
				NW 11%		
				W 33%		
				SW 8%		
				Plain 12%		
Velký Týnec	08.04.	254 m	2.3°	N 16%	2000	10.04.
				NW 45%		
				W 27%		
				SW 11%		
				Plain 1%		
Čechy	10.04.	222 m	0.8°	E 6%	1997	19.04.
				NE 10%		
				N 11%		
				NW 1%		
				SW 1%		
				Plain 72%		



ACKNOWLEDGEMENT:

The work was supported by Czech Ministry of Agriculture within the Research project no. QH 81049; "Integrated system of hop growing."