

Comparson of localsystemic and systemic Late Blight Fungicides

Product	Active ingredient	Application-rate/ha	MoA	Effect on Leaf blighth	Effect on stem blighth	Effect on new growth	Alternaria	Persistence	Kurative
localsystemic and systemic fungicides against late blight (<i>Phytophthora infestans</i>)									
Banjo forte	Dimethomorph 200 + Fluazinam 200 ¹⁾	1	H5 C5	☐	☐	○	☐	☐	☐
Carial Flex	Mandipropamid 250 + Cymoxanil 180	0.6	H5 U	☐	☐	○	○	●	☐
Curzate 60 WG	Cymoxanil 600	0.2	U	☐	☐	○	○	☐	☐
Infinito	Propamocarb 625 + Fluopicolide 62,5	1.6	F4 B5	☐	☐	☐	○	●	☐
Terminus Extra	Fluazinam 300 ¹⁾ + Cymoxanil 200	0.6	C5 U	☐	☐	○	○	☐	☐
Presidium	Dimethomorph 180 + Zoxamide 180	1	H5 B3	☐	☐	○	☐	☐	☐
Reboot	Cymoxanil 330 + Zoxamide 330	0.45	U B3	☐	☐	○	☐	☐	☐
Revus	Mandipropamid 250	0.6	H5	☐	☐	○	○	●	☐
Revus Top	Mandipropamid 250 + Difenconazol 250	0.6	H5 G1	☐	☐	○	☐	●	☐
Simpro	Propamocarb 400 + Cymoxanil 50	2.5	F4 U	☐	☐	☐	○	☐	☐
Voyager	Valifenalate 150 + Fluazinam 200 ¹⁾	1	H5 C5	☐	☐	○	☐	●	☐
Zorvec Entecta	Oxathiapiprolin 48 + Amisulbrom 240	0.25	F9 C4	●	☐	☐	○	●	☐

Rating: ○ = No effect; ☐ = very low; ☐ = low; ☐ = low to mediuml; ● = medium; ● = medium to gooe; ● = good

last update: march 2024

1) The classification assumes a Phytophthora population sensitive to fluazinam or cyazofamid.

