

FIGURE 1. EFFECT OF ISO-ALPHA-ACIDS ON CELL MEMBRANES.

METHODS

- iso- α -acids added to modified MRS broth, pH range 3.6 – 6.4
- inoculated with beer spoilage *Lactobacillus brevis*
- incubation (in dark) at 25°C for 48 hours
- Minimum Inhibitory Concentration (MIC) of each compound determined by measuring absorbance at 560 nm
- hydrophobicity determined using reverse-phase HPLC (Hughes et al., 1996)

The antibacterial activities of 12 iso- α -acids were investigated:

- cis*- & *trans*- isomers of 3 naturally occurring iso- α -acids
- 5 chemically reduced iso- α -acids
- reduced iso- α -acid mixture

i.e.

TIC	<i>trans</i> -isocohumulone
CIC	<i>cis</i> -isocohumulone
TIH	<i>trans</i> -isohumulone
TIA	<i>trans</i> -isoadhumulone
CIH	<i>cis</i> -isohumulone
CIA	<i>cis</i> -isoadhumulone
RIH	reduced iso- α -acids
DH1	dihydroiso- α -acid 1
DH2	dihydroiso- α -acid 2
TTIC	<i>trans</i> -tetrahydroisocohumulone
HIC	hexahydroisocohumulone
HIH	hexahydroisohumulone

INTRODUCTION

Hops are used to impart bitterness and aromatic flavours to beer, however certain compounds also possess antimicrobial properties. The iso- α -acids in particular possess strong antibacterial action against Gram positive bacteria (Simpson & Smith, 1992).

Iso- α -acids act as mobile carrier type ionophores, causing breakdown of the trans-membrane proton gradient of susceptible cells (Simpson, 1993). The cell is therefore unable to take up nutrients (Figure 1).




FIGURE 2. ISO-ALPHA-ACIDS STUDIED

Increased hydrophobicity leads to a greater antimicrobial activity (Figure 3). The more hydrophobic, reduced iso- α -acids are more antimicrobial than their naturally occurring analogues, and the degree of reduction is important (Figure 4). Increased hydrophobicity results in increased lipophilicity. This would render a compound more prone to interaction with the cell membrane, which could explain the observed effects.

References

- Hughes, P.S., Wilde, P.J. & Meneer, I.D., *Eur. Brew. Conv. Beer Foam Sub-Group Meeting*, 1996, 74-91
 Shimwell, J.L., *J. Inst. Brew.*, 1937, **43**, 191-195
 Simpson, W.J. & Smith, A.R.W., *J. Appl. Bacteriol.*, 1992, **72**, 327-334
 Simpson, W.J., *J. Inst. Brew.*, 1993, **139**, 1041-1045




FIGURE 3. EFFECT OF pH AND HYDROPHOBICITY ON ANTIMICROBIAL ACTIVITY (n.b. the lower the MIC value, the higher the antimicrobial activity)

Figure 3 also shows that antimicrobial activity is higher at lower pH (first noted by Shimwell, 1937). It is the undissociated forms of the iso- α -acids that possess antimicrobial properties. As they are weakly acidic, at lower pH there will be a greater concentration of the undissociated iso- α -acids.




FIGURE 4. DEGREE OF HYDROPHOBICITY OF HOP COMPOUNDS.

CONCLUSIONS

- reduced hydrophobicity & lower pH \uparrow increased antimicrobial activity.
- chemically reduced iso- α -acids \uparrow more antimicrobial.
- reduced hop compounds often used to prevent 'lightstruck' flavour.
- for sterile filtered, unpasteurised beers that are prone to microbiological problems, the addition of reduced hop compounds \uparrow additional benefit of improved microbiological stability.

Acknowledgements

Sheila Lee for technical assistance, Louise Bolshaw for purification of hop compounds and Paul Hughes for hydrophobicity data