

# HOP VIROIDS, HLVD AND HSVd BIOLISTIC INFECTION AND QUASIFORMS ANALYSIS

Matoušek J., Orctová L., Patzak, J. and Svoboda P. Institute of Plant Molecular Biology, Acad. Sci. CR, České Budějovice, Hop Institute GmbH, Žatec, Czech Republic. [jmat@umbr.cas.cz](mailto:jmat@umbr.cas.cz)

<http://www.umbr.cas.cz/804/www/>



Acknowledgements: W. Bardi, M. Hrdá and M. Mikušová (ITMB, AS CR) for their excellent technical assistance. This work was supported by NAIN7-MD-Q113 and GA AVCR 5503/11/4.

Molecular analysis of two hop viroids, HSVd and HLVD is performed with respect to their variability and ability to infect various weed species potentially important for viroid spreading and microevolution (see Figs on the side of the poster). While HLVD was referred to have the host range limited to hop, HSVd forms very invasive group with many molecular variants or quaspecies and ability to infect various plant species. Recent analyses (Matoušek J. e.a.: Virology 287:349, 2001) show, however, that HLVD forms quaspecies, too, providing its replication under temperature stress. Moreover, it was found that certain thermotomats of HLVD can undergo microevolution in *Solanaceae* species forming "low level" populations (Matoušek J.: Biol. Plant, 46: 607, 2003). Hence, in principle, under certain conditions both these viroids could have some potential to propagate in various weed species, perhaps forming "low level" populations representing latent danger for various hop genotypes.



Fig. 3






Fig. 4

800

HLVD infectious construct: cDNA with duplication of Pst I cleavage site, amplification with primers as shown in Fig. 1 below



Eco R1 restriction site was used to prepare infectious cDNAs of HSVd (Fig. 2)



Galinsoga ciliata (Raf.) Blake



Epilobium ciliatum Schreb



Urtica sp.



Artemisia vulgaris L.



Galium aparine L.

In the present work biolistic methods for viroid transfer in infectious cDNA (Fig. 1 and 2) and RNA (12-20 % PEG fraction as described by Matoušek e. a. J. Plant Physiol. 133:401, 1988) forms were developed. It was found that a minute amount of viroid RNA is sufficient to promote viroid infection by this approach, which is about 1000x more efficient than classical inoculation using carborundum method. HSVd variants that were collected from grapevines surrounding hoggardens (Matoušek e. a.: Plant Soil Environ. 49:168, 2003) were easily transmissible to hop, in which it promotes severe symptoms and developmental distortions (Fig. 3). Using TGGE and cDNA heteroduplex analysis, we found that at least 70 % of grapevine samples from locations close to hoggardens in Northern Bohemia [Žatec and Ústíek growing areas], but also in Slovenia [Zalec growing area] were infected with HSVd forming populations containing quaspecies (Fig. 4). Particular variant of HSVd corresponding to AC E01844 was experimentally transmissible from these samples to Czech hop Osvald's 72 also by classical mechanical inoculation. First attempts to inoculate various weed species led to finding that the strongest hybridisation signals are detectable in some biolistically inoculated members of *Compositae* family, such as *Galinsoga ciliata* (designated by the asterisk). In some plants like in *Urtica* sp. extremely low viroid concentration led to storn pathogenesis effect. Analysis of HLVD and HSVd propagation in approx. 15 weed species collected in hoggarden agrobiotype is in progress.



Erysimum cheiranthoides L.



Capsella bursa-pastoris (L.)



Sonchus oleraceus L.



Taraxacum officinale L.




Senecio vulgaris L.

## Conclusions

1. Highly efficient method of biolistic inoculation of hop viroids was developed.
2. Biolistic transfer of natural quasiforms of HSVd/HLVD to Osvald's 72 clone resulted in strong viroid infection and HSVd symptoms development on this hop.
3. Analysis of viroid infections after biolistic inoculation into various weed plant species was performed. Strong hybridization signals were detected in samples from *G. ciliata* and in some other plants mainly from *Compositae* family. Complex RT-PCR analysis is in progress.

Symptoms on *G. ciliata* leaves



HLVD calibration from hop



Symptoms on Urtica



RT-PCR detector of HLVD U.U, H.H, 0

Galinsoga ciliata

Artemisia vulgaris

Urtica dioica L.

Achillea millefolium

Senecio vulgaris